



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

“The three points in which any line cuts the sides of a triangle, and the projections, from any point in the plane, of the vertices of the triangle on the same line are six points in involution.”

The fact that this proposition has thus been derived from another as its reciprocal, will not at all interfere with the deduction in the ordinary manner of the reciprocal theorem concerning six rays in involution joining any point to the vertices of a complete quadrilateral.

On the other hand, as no use whatever has been made of harmonic properties to obtain the theorem, these may all be deduced as special applications, by drawing the transversal AX through two of the points in which sides of the triangle MNP meet the lines joining Q to the opposite vertices, thus making these points foci of the involution.

Or, should it be preferred to proceed more nearly in the usual course, a definition of the punctual distance, equivalent to the one already stated, may be given in terms of angular functions. Thus, in the figure, the punctual distance between NC' and NB is

$$\frac{\sin NAX \sin BNC'}{NN \sin ANB \sin ANC'};$$

and it is easily shown that the ratio in which the punctual distance of two out of three convergents is divided by the third is equal to the anharmonic ratio of the pencil formed by adding to the three convergents a line which joins the point of convergence to the origin.

A NEW AND USEFUL FORMULA FOR INTEGRATING CERTAIN DIFFERENTIALS.

BY PROF. J. W. NICHOLSON, A. M., BATON ROUGE, LA.

PROBLEM.—*To integrate $v^n du$ in terms of the descending powers of v .*
Let us assume

$$\int v^n du = yv^n + y_1 v^{n-1} + y_2 v^{n-2} + \dots + y_t v^{n-t} + \int z v^{n-t-1} dv. \quad (1)$$

Differentiating (1), and arranging with reference to v , we have

$$0 = - \frac{du}{dy} \Big| v^n + \frac{ny dv}{dy_1} \Big| v^{n-1} + (n-1) \frac{y_1 dv}{dy_2} \Big| v^{n-2} + \dots + (n-t) \frac{y_t dv}{z dv} \Big| v^{n-t-1} \quad (2)$$

Now since (2) is true for every value of v , according to the principle of indeterminate coefficients, we have

$$\left. \begin{aligned} dy &= du, \\ dy_1 &= -nydv, \\ dy_2 &= -(n-1)y_1dv, \\ dy_3 &= -(n-2)y_2dv, \\ &\dots\dots\dots \\ dy_n &= -(n-t+1)y_{t-1}dv, \\ z &= -(n-t)y_t. \end{aligned} \right\} \quad (3)$$

Integrating equations (3), supposing the constant to be added = 0, and denoting

$$\begin{aligned} f dv f u dv &\text{ by } f^2 u dv^2, \\ f dv f dv f u dv &\text{ by } f^3 u dv^3, \\ \&c. \quad \&c. \quad \&c., \end{aligned}$$

we have

$$\left. \begin{aligned} y &= u \\ y_1 &= (-1)^1 n f u dv \\ y_2 &= (-1)^2 n(n-1) f^2 u dv^2 \\ y_3 &= (-1)^3 n(n-1)(n-2) f^3 u dv^3 \\ &\dots\dots\dots \\ y_t &= (-1)^t n(n-1) \dots (n-t+1) f^t u dv^t \\ z &= (-1)^{t+1} n(n-1) \dots (n-t) f^t u dv^t. \end{aligned} \right\} \quad (4)$$

Substituting in (1), we have

$$\begin{aligned} f v^n du &= u v^n - n v^{n-1} f u dv + n(n-1) v^{n-2} f^2 u dv^2 - n(n-1)(n-2) v^{n-3} f^3 u dv^3 \\ &+ \dots (-1)^t n(n-1) \dots (n-t+1) v^{n-t} f^t u dv^t + \dots (-1)^{t+1} n(n-1) \dots \\ &\quad (n-t) f v^{n-t-1} dv f^t u dv^t. \end{aligned} \quad (5)$$

If $t = n$ (5) becomes

$$\begin{aligned} f v^n du &= u v^n - n v^{n-1} f u dv + n(n-1) v^{n-2} f^2 u dv^2 \dots (-1)^n n(n-1) \dots \\ &\quad \dots (3) (2) (1) f^n u dv^n. \end{aligned} \quad (6)$$

Of course (6) is not so useful and general as (5), but the want of space will scarcely allow us to indicate the utility of the former in the applications of the formula which we purpose make to the integration of a few important differentials.

Ex. 1. $\int x^m \log x dx = ?$

Make $u = x^{m-1} \div (m+1)$, and $v = \log x$. Hence,

$$\begin{aligned} u dv &= \frac{x^m dx}{m+1}, \quad \int u dv = \frac{x^{m+1}}{(m+1)^2}; \\ dv \int u dv &= \frac{x^m dx}{(m+1)^2}, \quad \int^2 u dv^2 = \frac{x^{m+1}}{(m+1)^3}. \end{aligned}$$

Similarly,

$$\begin{aligned} \int^3 u dv^3 &= \frac{x^{m+1}}{(m+1)^4}, \\ &\dots\dots\dots \\ \int^n u dv^n &= \frac{x^{m+1}}{(m+1)^n}. \end{aligned}$$

Substituting in (6), we have

$$\int x^m \log^n x dx = \frac{x^{m+1}}{m+1} \left[\log^n x - \frac{n \log^{n-1} x}{m+1} + \frac{n(n-1) \log^{n-2} x}{(m+1)^2} \dots \right. \\ \left. (-1)^n \frac{n(n-1) \dots (3)(2)(1)}{(m+1)^n} \right]. \quad (7)$$

If in (7) $m = 0$, and we take the integral between the limits $x = 0$ and $x = 1$, we have

$$\int_0^1 \log^n x dx = (-1)^n n(n-1)(n-2) \dots 3 \cdot 2 \cdot 1.$$

Ex. 2. $\int a^x x^n dx = ?$

Make $u = a^x \div (\log a)$, and $v = x$. Hence,

$$\begin{aligned} \int u dv &= \frac{a^x}{\log^2 a}, \\ \int u dv^2 &= \frac{a^x}{\log^3 a}, \\ &\dots \dots \dots \\ \int u dv^n &= \frac{a^x}{\log^{n+1} a}. \end{aligned}$$

Substituting in (6),

$$\int a^x x^n dx = \frac{a^x}{\log a} \left[x^n - \frac{nx^{n-1}}{\log a} + \frac{n(n-1)x^{n-2}}{\log^2 a} \dots (-1)^n \frac{n(n-1) \dots 3 \cdot 2 \cdot 1}{\log^n a} \right].$$

In a similar manner we may integrate a number of other important differentials, such as

$$\sin^m x \cos^n x dx, \quad (a + bx^n)^p x^n dx, \quad \&c.$$

SOLUTION OF PROB. 334, BY W. E. HEAL.—(See pp. 60, 98, Vol. VIII)

Let the given ellipse be

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

The locus of intersection of tangents which intersect at a constant angle, φ , is

$$[(x^2 + y^2) - (a^2 + b^2)]^2 \tan^2 \varphi = 4(b^2 x^2 + a^2 y^2 - a^2 b^2). \quad (A)$$

(Salmon's Conics, 5th edition, page 161.)

The polar of every point on this curve touches the required curve and conversely the polar of every point on the required curve touches the curve (A).

Let (x, y) be the coordinates of any point on the required curve.

The polar of this point is

$$\frac{xx'}{a^2} + \frac{yy'}{b^2} = 1. \quad (B)$$